SIDE B



- If you must cancel your appointment, we require 48 hours notice. (Products are purchased specifically for your procedure) Please call 613 969-7400 Ext. 2494
- Check in at Nuclear Medicine Take the Staff
 Elevators in the Quinte Wing to the Basement Level and follow the arrows.

Appointment Date and Time:

 Your test can take approximately 4.0 hours

NOTE: Your test may be rebooked if instructions are not followed.

Instruction for Myocardial Perfusion (MIBI) Test

Dipyridamole

(After consultation with your physician, and unless otherwise directed, follow these instructions)

36 hours before your appointment:

X STOP The following medications;
Aminophylline, Uniphyl, Theodur, Trental

24 hours before your appointment:

- X STOP All Beta Blockers, such as;
 Metoprolol, Bisoprolol, Atenolol, Carvedilol,
 Monocor, Lopresor, Sotalol, Propanolol
- X STOP All Calcium Antagonists, such as; Norvasc, Cardizem, Diltiazem, Adalat, Amlodipine, Tiazac, Felodipine, Nifedipine, Verapamil, Plendil
- X STOP Long Acting Nitrates, such as; IsoBid, Isordil, Isorbide dinitrate, Imdur Sorbitrate Dilatrate-SR
- **X STOP** Aggrenox
- X STOP Viagra, or Cialis
- X STOP Gravol, Fiorinal, Tylenol 1, 2, or 3 or any meds containing caffeine(ask Pharmacist)

12 hours before your appointment:

X STOP Nitro Patch, but continue to use Nitro Spray or Nitro pills as needed.



What can I eat and drink?

24 hours(1 day) before appointment; Eat your usual meals, **EXCEPT;**

X STOP All types of coffee and tea, including decaf and herbal

X STOP All types of chocolate, including hot chocolate, and chocolate milk

X STOP All soft drinks, pop, energy drinks

ON THE DAY OF YOUR TEST:

Bring all of your medications in their original containers.

You must bring food or money to purchase food.

Fast 4 hours, only water permitted.

Wear comfortable clothes and shoes. (NO SANDALS OR HEELS.)

No Smoking on the morning of test

Make sure you have followed the instructions for the procedure.